

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/48, 9/70	A1	(11) International Publication Number: WO 96/40139 (43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: PCT/US96/09692 (22) International Filing Date: 7 June 1996 (07.06.96) (30) Priority Data: 08/473,631 7 June 1995 (07.06.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/473,631 (CIP) Filed on 7 June 1995 (07.06.95) (71) Applicant (for all designated States except US): ALZA CORPORATION [US/US]; 950 Page Mill Road, P.O. Box 10950, Palo Alto, CA 94303-0802 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): YUM, Su II [US/US]; 1021 Runnymede Court, Los Altos, Ca 94022 (US). NELSON, Melinda, K. [US/US]; 1127 Hollenbeck Road, Sunnyvale, CA 94087 (US). CAMPBELL, Patricia, S. [US/US]; 1410 Middlefield Road, Palo Alto, CA 94301 (US). (74) Agent: RAFA, Michael, J.; Alza Corporation, 950 Page Mill Road, P.O. Box 10950, Palo Alto, CA 94303-0802 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NOVEL FORMULATIONS FOR TRANSDERMAL DELIVERY OF PERGOLIDE		
(57) Abstract <p>Composition of matter for application to a body surface or membrane to administer pergolide by permeation through the body surface or membrane, the composition comprising pergolide to be administered, at a therapeutically effective rate, alone or in combination with a permeation enhancer or mixture. Also disclosed are drug delivery devices containing the pergolide or pergolide and enhancer composition and methods for the transdermal administration of the pergolide and pergolide/enhancer composition.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**NOVEL FORMULATIONS FOR
TRANSDERMAL DELIVERY OF PERGOLIDE**

FIELD OF INVENTION

5

This invention relates to the safe and efficacious transdermal administration of pergolide for, among other things, the treatment of Parkinson's Disease. More particularly, the invention relates to novel methods, compositions, and devices for administering pergolide to a subject through a body surface or membrane over a sustained time period. A preferred embodiment is directed to the transdermal co-administration of a pharmaceutically acceptable salt of pergolide in combination with a permeation enhancer.

15

BACKGROUND OF THE INVENTION

Pergolide, 8-[(methylthio)methyl]-6-propylergoline, a compound based on the ergoline ring system, is reported to be a dopaminergic agonist that also decreases plasma prolactin concentrations. When used for treating Parkinson's Disease, pergolide is used as a conjunctive therapy with levodopa.

US Patent No. 4,166,182, incorporated herein in its entirety by reference, describes the preparation of pergolide and its oral or parenteral administration as a prolactin inhibitor and in the treatment of Parkinson's Disease.

German patent application DE 4240798, incorporated herein its entirety by reference, describes a pharmaceutical composition containing ergot derivatives, including pergolide, for protection of nerves. The composition may be delivered orally, sublingually, parenterally, percutaneously or nasally.

US Patent No. 4,797,405 incorporated herein in its entirety by reference, discusses stabilized pergolide oral compositions that demonstrate reduced decomposition when exposed to light.

The dopaminergic agonist effect of pergolide has resulted in its use in a variety of treatments, in addition to the treatment of Parkinson's Disease. For example, US Patent No. 4,800,204, incorporated herein in its entirety by reference, discusses a method of controlling tobacco use by orally or parenterally administering a direct dopamine receptor agonist such as pergolide.

US Patent No. 4,935,429, incorporated herein in its entirety by reference, discusses a method of treating psychostimulant abuse by orally or parenterally administering a dopamine agonist such as pergolide.

US Patent No. 5,063,234, incorporated herein in its entirety by reference, discusses a method of inhibiting bone demineralization by administering, preferably orally, an ergot derivative, such as pergolide.

The oral administration of pergolide in the treatment of Parkinson's Disease is initiated with 0.05 mg/day dosage for the first 2 days. The dosage is then gradually increased by 0.1 or 0.15 mg/day every third day over the next 12 days of therapy. The dosage may then be increased by 0.25 mg/day every third day until an optimum therapeutic dosage is achieved at a range of about 1.5 to 8.0 mg/day. Generally, the daily dose is divided into three oral doses. The side effects of oral administration include, but are not limited to nausea, vomiting, dizziness and orthostatic hypotension.

The transdermal route of parenteral delivery of drugs and other biologically active agents ("agents") has been proposed for a wide variety of systemically acting and locally acting agents on either a rate-controlled or non-rate-controlled basis and is described in numerous technical publications such as the following: US Patents 3,598,122; 3,598,123; 3,731,683; 3,797,494; 4,031,894; 4,201,211; 4,286,592; 4,314,557; 4,379,454; 4,435,180; 4,559,222; 4,568,343; 4,573,995; 4,588,580; 4,645,502; 4,704,282; 4,788,062; 4,816,258; 4,849,226; 4,908,027; 4,943,435; and

5,004,610, the disclosures of which are incorporated in their entirety herein by reference. The transdermal administration of a related compound, lisuride, for treating Parkinson's Disease, is disclosed in US Patents 5,252,335 and 5,229,129, the disclosures of which are incorporated in their entirety herein by reference.

When first investigated in depth in the late 1960's, the transdermal route of administration appeared to offer many advantages, particularly with respect to agents that had short half lives and therefore required frequent, repeated dosing or were subject to a high degree of first-pass metabolism by the liver when orally administered. The peaks and valleys in blood concentration resulting from frequent periodic doses of short half-life agents would be eliminated and replaced by substantially constant plasma concentration. This would not only improve individual compliance but also would eliminate the alternating periods of high side-effects and ineffective blood concentrations associated with period dosing. Administering the agent through the skin directly into the blood stream would also eliminate first-pass metabolism of orally administered agents.

It was initially assumed, theoretically, that any short half-life agent of high potency and skin permeability would be suitable for safe and effective transdermal administration. This assumption, however, has not been proven true.

The failure of the transdermal route to fulfill the initial expectations of its potential as an administrative portal was primarily due to the incredible variety of properties with which nature has endowed the skin to permit it to perform its function as the primary barrier to prevent the ingress of foreign substances into the body. See Transdermal Drug Delivery: Problems and Possibilities, B. M. Knepp, et al, CRC Critical Reviews and Therapeutic Drug Carrier Systems, Vol. 4, Issue 1 (1987).

Thus, the transdermal route of administration, rather than being available to every short half-life agent of high potency and skin permeability, was found to be available only to those few agents that possess the proper

combination of a host of characteristics, most of which are unpredictable, required to render the agent suitable for safe and effective transdermal administration.

The most significant of these characteristics are the following:

- 5 1. Skin Permeability. The permeability of the skin to the agent must be sufficiently high so that the agent can be administered at a therapeutically effective rate through an area of skin no greater than about 200 cm² and preferably no greater than 50 cm². The person-to-person variation in skin permeability at similar sites should also be considered.
- 10 2. Skin Binding. The skin beneath the transdermal delivery device has the capability of creating a skin depot of drug by absorbing, adsorbing, or binding a certain amount of agent. The amount of agent so bound must be supplied to the skin before the agent can be delivered into the blood stream at steady, therapeutically effective rates. If large amounts of the agent are
15 bound in the skin, significant delays in the onset of therapeutic effect ("lag time") will be observed together with corresponding delays and termination of effect upon removal of the device. The potential also exists for toxic quantities of potent agents to be contained within the skin beneath the device. Skin binding is not related to skin permeability. Agents that are
20 highly permeable may also be highly bound causing a lag time sufficiently long as to render them unsuitable for their intended use.
3. Irritation. The skin reacts to many topically applied substances, particularly those maintained under occlusion, by blistering or reddening accompanied by unpleasant burning, itching, and stinging sensations.
25 Animal models are used to screen for irritation. Animal models, however, often produce both false positives and false negatives. There is also a wide interpersonal variation in susceptibility to irritation. An agent must be minimally irritating in a large percentage of the potential individual population in order to be suitable for safe and effective transdermal administration.
- 30 4. Sensitization. Sensitization is an allergic reaction which is induced when an agent is first applied to the skin and is elicited upon

continued exposure which may occur immediately or after a long period of seemingly harmless exposure.

The sensitization may be local, elicited by topical exposure, which manifests itself as contact dermatitis accompanied by blistering, itching, reddening and burning at the site of application. More seriously, the sensitization may be systemic, elicited by topical application but manifesting itself by more general allergic reactions at sites other than the site of application. Most seriously, the systemic sensitization may be elicited by oral or intravenous administration of the drug. If the latter occurs, the individual will be unable to take the drug by any route of administration.

Animal models are used to screen for sensitization. Animal models, however, produce both false positives and false negatives. There is also a wide variation in the allergic reaction between individuals as well as between sexes, races and skin types. It is obvious that a useful transdermal agent must be minimally sensitizing in a large percentage of the potential individual population.

5. Pharmacokinetic Properties. The half-life of an agent is the time after administration that half of the amount administered has been eliminated from the body. Because blood concentrations of continuously administered agents will continue to increase for approximately six half-lives before steady-state constant blood concentrations are achieved, an agent must have a relatively short half-life to be suitable for continuous transdermal administration. The transdermal half-lives of most agents have not been determined. When half-lives of agents determined from intravenous administration are compared with half-lives determined from transdermal administration, the transdermal half-lives are generally longer but there can be wide variation in half-life between individuals based upon factors such as age, sex, health, and body type.

6. Pharmacodynamic Properties. Constant blood levels may not produce the desired therapeutic effects. For example, a therapeutic effect may only be observed at peak blood concentration obtained from bolus

dosing but the peak concentration cannot be maintained because of side effects associated therewith. Also, continuous administration of many agents produces tolerance that may require either some agent-free interval or continually increasing and therefore potentially hazardous doses of the agent.

7. Potency. Although a certain degree of potency is required for transdermally administered agent to be effective, it is also possible for an agent to be too potent. As potency increases, lower blood concentrations are required and much smaller quantities are administered. Because of normal inter-individual variations and skin permeability, it may not be possible to precisely control whether a individual is receiving 1 $\mu\text{g/hr}$ or 2 $\mu\text{g/hr}$, for example. For a highly potent agent, a 1 $\mu\text{g/hr}$ administration may be totally ineffective and a 2 $\mu\text{g/hr}$ rate fatal. Thus, the therapeutic index of an agent, which is the ratio of toxic blood concentration to the therapeutic blood concentration, becomes extremely significant. A highly potent agent should also have a relatively wide therapeutic window in order to be suitable for transdermal administration.

8. Metabolism. One of the perceived advantages of transdermal administration was that it avoided the "first-pass" metabolism of the agent by the liver that is associated with oral administration. It has now been recognized, however, that the skin is also a large metabolizing organ in the body for some drugs. Thus, although first-pass metabolism that occurs after an orally administered agent enters the blood stream can be avoided, skin metabolism, which occurs before the agent enters the bloodstream, cannot be avoided. Skin metabolism is capable of creating metabolites that are inert, irritating, toxic, or comparable in biological activity to that of the agent. An agent, to be suitable for transdermal administration, must have the metabolic properties that are consistent with its therapeutic use on continuous administration.

The above summarizes the primary characteristics that effect suitability of an agent for transdermal administration that have been recognized to date.

There are undoubtedly others, some of which have not yet been recognized, and, in order for an agent to be suitable for transdermal administration, it must possess the right combination of all these characteristics, a combination of which, as illustrated by the very few drugs that are now suitable for administration from transdermal delivery devices, is quite rare and unpredictable.

SUMMARY OF THE INVENTION

It is unexpected that pergolide would be delivered through the skin at meaningful therapeutic rates either as a base or salt because, as its chemical name 8-[(methylthiomethyl)-6-propylergoline monomethanesulfonate]] indicates, it has a complex chemical structure which does not lend itself to readily permeate through biological membranes such as the skin.

Furthermore, it is even more unexpected that transdermal dosage forms provided with a pharmaceutically acceptable salt form of pergolide would result in a greater pergolide flux through skin than dosage forms provided with pergolide free base.

Nonetheless, according to this invention, it has been discovered that pergolide can be safely and efficaciously administered transdermally to provide, among other things, treatment for Parkinson's Disease, with a reduced incidence of side effects and improved individual compliance. In addition, the present invention provides methods for the transdermal delivery of pergolide and delivery systems for effecting the same, which are suitable for the administration of pergolide continuously through a body surface or membrane to achieve and maintain therapeutic blood plasma levels of pergolide in an individual. A particularly advantageous aspect of this invention is the ability to maintain substantially constant blood plasma levels of pergolide in an individual over extended periods of time.

Surprisingly, the inventors have discovered that transdermal dosage forms provided with a pharmaceutically acceptable salt form of pergolide

result in a greater pergolide flux through skin than dosage forms provided with pergolide free base. This is contrary to findings of a substantial amount of the prior art wherein for a majority of agents which have been tested for transdermal administration, it is recognized in the art that the permeability of the skin to the unionized form of the drug is generally substantially greater than that of the ionized form. A preferred embodiment is therefore directed to providing the reservoir of a transdermal delivery device with a pharmaceutically acceptable salt of pergolide, preferably pergolide mesylate, together with a permeation enhancer.

10 As used herein, the term "transdermal" intends percutaneous and transmucosal administration, ie, passage of pergolide through intact unbroken skin or mucosal tissue into the systemic circulation.

As used herein, the term "pergolide" intends not only the basic form of pergolide but also pharmaceutically acceptable salt forms of pergolide.

15 As used herein the term "salt" intends, but is not limited to, pharmaceutically acceptable salts such as chlorides, acetates, sulfates, phosphates, and mesylates.

As used herein, the term "pergolide therapy" intends all medical conditions for which pergolide is or will be indicated, including, without limitation, as a psychic energizer and in the treatment of Parkinson's Disease, migraine, allergic responses, urticaria, hypertension, endometritis, and other conditions associated with dopaminergic agonists.

20 As used herein, the term "individual" intends a living mammal and includes, without limitation, humans and other primates, livestock and sports animals such as cattle, pigs and horses, and pets such as cats and dogs.

As used herein, the term "therapeutic blood plasma level" intends the level of pergolide in blood plasma that achieves a therapeutic effect and is typically within the range of about 100 pg/mL - 2000 pg/mL.

30 As used herein, the term "therapeutically effective rate" intends a rate of pergolide delivery effective to achieve therapeutic blood plasma levels of pergolide in an individual during the administration period.

As used herein, the phrase "sustained time period" or "administration period" intends at least about 8 hours and will typically intend a period in the range of about one to about seven days.

As used herein, the phrase "predetermined area of skin" intends a
5 defined area of intact unbroken skin or mucosal tissue. That area will usually be in the range of about 1 cm² to about 100 cm².

As used herein, the term "permeation enhancer" intends an agent or a mixture of agents which increases the permeability of the skin to pergolide.

As used herein, the term "permeation enhancement" intends an
10 increase in the permeability of skin to pergolide in the presence of a permeation enhancer as compared to permeability of skin to pergolide in the absence of a permeation enhancer.

The present invention relates to compositions, devices and methods for the transdermal administration of pergolide in the treatment of Parkinson's
15 Disease, among other things. According to the present invention, it has been found that pergolide may be safely and efficaciously administered transdermally through a body surface or membrane at a therapeutically effective rate for a predetermined, sustained time period in order to provide an effective therapeutic result. Another aspect of the present invention is
20 directed to the transdermal administration of pergolide together with a suitable permeation enhancer or mixture of enhancers.

The system of the invention comprises a carrier or matrix adapted to be placed in pergolide or pergolide- and permeation enhancer- transmitting relation to the selected skin or other body site. The carrier or matrix contains
25 sufficient amounts of pergolide to continuously administer pergolide to the site at a therapeutically effective rate during an administration period. In another embodiment, the carrier or matrix contains sufficient amounts of pergolide and a permeation enhancer to continuously coadminister pergolide at a therapeutically effective rate to the site together with the permeation
30 enhancer during the administration period.

The accompanying figures, which are incorporated in and constitute a part of this specification, illustrate embodiments of the invention and, together with the description serve to explain the principles of the invention.

5

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a cross-section through a schematic perspective view of one embodiment of a transdermal therapeutic system according to this invention prior to application to the skin.

10

Figure 2 is a cross-section view through another embodiment of this invention.

Figure 3 is a cross-section view through another embodiment of this invention.

15

Figure 4 is a cross-section view through another embodiment of this invention.

Figure 5 is a graph of the flux of pergolide base through human epidermis, in vitro, at 35°C, from various aqueous and non-aqueous donors.

Figure 6 is a graph of the flux of pergolide mesylate through human epidermis, in vitro, at 35°C, from various aqueous and non-aqueous donors.

20

Figure 7 is a graph of the cumulative release of pergolide base through human epidermis at 35°C, in vitro from an EVA matrix system with various permeation enhancers.

25

Figure 8 is a graph of the cumulative release of pergolide mesylate through human epidermis at 35°C, in vitro, from an EVA matrix system with various permeation enhancers.

Figure 9 is a graph of the cumulative release of pergolide base through human epidermis at 35°C in vitro, from an EVA matrix system with various permeation enhancers.

30

Figure 10 is a graph of the cumulative release of pergolide mesylate through human epidermis at 35°C, in vitro, from an EVA matrix system with various permeation enhancers.

Figure 11 is a graph of the cumulative release of pergolide base through human epidermis in vitro at 35°C, from an EtOH system having an EVA (18% vinyl acetate) rate controlling membrane with various permeation enhancers and a polyisobutylene/mineral oil based in-line contact adhesive preloaded with drug.

Figure 12 is a graph of the cumulative release of pergolide mesylate through human epidermis, in vitro, at 35°C, from an EtOH system having an EVA (18% vinyl acetate) rate controlling membrane with various permeation enhancers and a polyisobutylene / mineral oil based in-line contact adhesive preloaded with drug.

Figure 13 is a graph of the flux of pergolide base through human epidermis, in vitro, at 35°C, from an EtOH system having an EVA (28% vinyl acetate) rate controlling membrane with various permeation enhancers and a polyisobutylene/mineral oil based in-line contact adhesive preloaded with drug.

Figure 14 is a graph of the flux of pergolide mesylate through human epidermis, in vitro, at 35°C, from an EtOH system having an EVA (28% vinyl acetate) rate controlling membrane with various permeation enhancers and a polyisobutylene/mineral oil based in-line contact adhesive preloaded with drug.

Figure 15 is a graph of the flux of pergolide mesylate through human epidermis, in vitro, at 35°C, from various pergolide mesylate/GML/methyl laurate/EVA 40 systems with no adhesive.

Figure 16 is a graph of the flux of pergolide mesylate through human epidermis, in vitro, at 35°C, from various pergolide mesylate/GML/dodecyl acetate/EVA 40 systems with no adhesive.

Figure 17 is a graph of the flux of pergolide mesylate through human epidermis, in vitro, at 35°C, from various pergolide mesylate/GML/lauryl lactate/EVA 40 systems with no adhesive.

Figure 18 is a graph of the flux comparison of GML coenhancers methyl laurate, lauryl lactate, and dodecyl acetate in combination with mineral

oil through human epidermis, in vitro, at 35°C, from various pergolide mesylate/GML/coenhancer/mineral oil/EVA 40 systems with no adhesive.

DETAILED DESCRIPTION OF THE INVENTION

5

According to the present invention, it has been found that pergolide may be administered to the human body at a therapeutically effective rate via the transdermal route for the purpose of treating Parkinson's Disease, among other things, when administered alone or coadministered with a suitable permeation enhancer. Therapeutic blood plasma levels of 100 pg/mL to 2000 pg/mL can be achieved according to this invention. Representative in vitro skin fluxes of pergolide through human skin are in the range of 0.1 - 8.0 $\mu\text{g}/\text{cm}^2 \cdot \text{hr}$, depending on the form of pergolide (base or salt), the permeation enhancer, and whether an in-line contact adhesive was present in the path of drug flow. According to this invention, it is preferable to administer pergolide at a flux of at least about 1.0 $\mu\text{g}/\text{cm}^2 \cdot \text{hr}$.

It is estimated that therapeutic blood plasma levels can be achieved within approximately 5 to 10 hours following application of the first patch. When a subsequent system is applied, a lag period of no pergolide delivery is not experienced due to the presence of a skin depot of pergolide remaining from the prior system, thus pergolide is continuously administered throughout subsequent system applications in order to maintain therapeutic blood plasma levels of pergolide over a sustained time period. The system is easily adapted for shorter or longer duration treatments, but generally 72 hours is the preferred duration for a single treatment.

The desired pergolide administration rate may be achieved by increasing or decreasing the surface area of the transdermal delivery device without effecting the flux. For example, for a pergolide skin flux of 1.1 $\mu\text{g}/\text{cm}^2 \cdot \text{hr}$, a patch having a surface area of about 60 cm^2 would deliver approximately 1.6 mg of pergolide over a 24 hour period.

The pergolide administration rate may also be increased by increasing the flux of pergolide through skin by the use of permeation enhancers which include, but are not limited to: monoglycerides such as glycerol monolaurate, glycerol monooleate or glycerol monolinoleate, lactate esters such as lauryl lactate, methyl laurate, caproyl lactic acid, lauramide diethanolamine (LDEA), dimethyl lauramide, polyethylene glycol-4 lauryl ether (Laureth-4), and ethanol, alone or in combinations of one or more. An embodiment of this invention also relates to codelivery of at least one of the permeation enhancers mentioned above to aid in the transdermal delivery of pergolide.

The present inventors also found that certain adhesives were preferred as the in-line contact adhesive when one was used in a therapeutic transdermal pergolide system. More particularly, it was found that systems using polyisobutylene adhesives as the in-line contact adhesive resulted in greater flux of pergolide through skin than when other adhesives, such as acrylate adhesives, were used.

Therefore, the present invention, in one embodiment, is directed to a composition of matter for administration to a body surface or membrane to deliver pergolide by permeation through the body surface or membrane at a therapeutically effective rate, wherein the composition comprises an amount of pergolide in a carrier effective to permit sustained release of pergolide at a therapeutically effective rate over an administration period of at least about 8 hours, wherein said therapeutically effective rate of pergolide release achieves therapeutic blood plasma levels of pergolide in an individual.

Another embodiment is directed to a composition of matter for administration to a body surface or membrane to deliver pergolide by permeation through the body surface or membrane at a therapeutically effective rate, wherein the composition comprises, in combination:

- (a) a therapeutically effective amount of pergolide; and
- (b) a permeation-enhancing amount of a permeation enhancer.

The drug may be present in the composition in an amount ranging from 0.1 to 20% by weight.

This invention finds particular usefulness in administering pergolide across skin. It is also useful, however, in administering pergolide across mucosa. According to our invention, pergolide with or without a permeation enhancer are placed in drug or drug and permeation enhancer- transmitting relationship to an appropriate body surface, preferably in a pharmaceutically acceptable carrier thereof, and maintained in place for the desired administration period.

The pergolide and the permeation enhancer, if used, are typically dispersed within a physiologically compatible matrix or carrier, as more fully described below, which may be applied directly to the body as an ointment, gel, cream, suppository or sublingual or buccal tablet. When used in the form of a liquid, ointment, lotion, cream or gel applied directly to the skin, it is preferable, although not required, to occlude the site of administration. Such compositions can also contain other permeation enhancers, stabilizers, dyes, diluents, pigments, vehicles, inert fillers, excipients, gelling agents, vasoconstrictors, and other components of topical compositions as are known to the art.

In other embodiments, the pergolide with or without the permeation enhancer would be administered from a transdermal device as more fully described below. Examples of suitable transdermal delivery devices are illustrated in Figs. 1 - 4. In the figures, the same reference numbers are used throughout the different figures to designate the same or similar components.

The figures are not drawn to scale.

Referring now to Figure 1, a preferred embodiment of a transdermal therapeutic system according to this invention comprises transdermal delivery device 10 comprising a reservoir 12, preferably in the form of a matrix containing pergolide and a permeation enhancer, if used, dispersed therein. Reservoir 12 is sandwiched between a backing 14 and an in-line contact adhesive layer 16. The device 10 adheres to the surface of the skin 18 by

means of the adhesive layer 16. The adhesive layer 16 may optionally contain the permeation enhancer and/or pergolide. A strippable release liner (not shown in FIG. 1) is normally provided along the exposed surface of adhesive layer 16 and is removed prior to application of device 10 to the skin

5 18. Optionally, a rate-controlling membrane (not shown) may be present between the reservoir 12 and the adhesive layer 16. Although the preferred embodiments of this invention utilize an in-line adhesive as is shown in Figure 1, other means for maintaining the system on the skin can be employed. Such means include a peripheral ring of adhesive outside the

10 path of the drug from the system to the skin or the use of other fastening means such as buckles, belts, and elastic arm bands.

Alternatively, as shown in FIG. 2, transdermal therapeutic device 20 may be attached to the skin or mucosa of a individual by means of an adhesive overlay 22. Device 20 is comprised of reservoir 12, preferably in

15 the form of a matrix containing pergolide and a permeation enhancer, if used, dispersed therein. A backing layer 14 is provided adjacent one surface of reservoir 12. Adhesive overlay 22 maintains the device on the skin and may be fabricated together with, or provided separately from, the remaining elements of the device. With certain formulations, the adhesive overlay 22

20 may be preferable to the in-line contact adhesive 16 as shown in FIG. 1. Backing layer 14 is preferably slightly larger than reservoir 12, and in this manner prevents the materials in reservoir 12 from adversely interacting with the adhesive in overlay 22. Optionally, a rate-controlling membrane (not shown in FIG. 2) may be provided on the skin-proximal side of reservoir 12.

25 A strippable release liner 24 is also provided with device 20 and is removed just prior to application of device 20 to the skin.

In FIG. 3, transdermal delivery device 30 comprises a pergolide and permeation enhancer reservoir ("pergolide reservoir") 12 substantially as described with respect to FIG. 1. Permeation enhancer reservoir ("enhancer reservoir") 26 comprises the permeation enhancer dispersed throughout and

30 contains pergolide at or below saturation, when in equilibrium with the first

reservoir. Enhancer reservoir 26 is preferably made from substantially the same matrix as is used to form pergolide reservoir 12. A rate-controlling membrane 28 for controlling the release rate of the permeation enhancer from enhancer reservoir 26 to pergolide reservoir 12 is placed between the two reservoirs. A rate-controlling membrane (not shown in FIG. 3) for controlling the release rate of the enhancer from pergolide reservoir 12 to the skin may also optionally be utilized and would be present between adhesive layer 16 and reservoir 12.

The rate-controlling membrane may be fabricated from permeable, semipermeable or microporous materials which are known in the art to control the rate of agents into and out of delivery devices and having a permeability to the permeation enhancer lower than that of drug reservoir 12. Suitable materials include, but are not limited to, polyethylene, polyvinyl acetate, ethylene n-butyl acetate and ethylene vinyl acetate copolymers.

Superimposed over the permeation enhancer reservoir 26 of device 30 is a backing 14. On the skin-proximal side of reservoir 12 are an adhesive layer 16 and a strippable liner 24 which would be removed prior to application of the device 30 to the skin.

In the embodiments of FIGS. 1, 2 and 3, the carrier or matrix material of the reservoirs has sufficient viscosity to maintain its shape without oozing or flowing. If, however, the matrix or carrier is a low-viscosity flowable material such as a liquid or a gel, the composition can be fully enclosed in a pouch or pocket, as known to the art from US Pat. No. 4,379,454 (noted above), for example, and as illustrated in FIG. 4. Device 40 shown in FIG. 4 comprises a backing member 14 which serves as a protective cover for the device, imparts structural support, and substantially keeps components in device 40 from escaping the device. Device 40 also includes reservoir 12, which contains the pergolide with or without a permeation enhancer, and bears on its surface distant from backing member 14, a rate-controlling membrane 28 for controlling the release of pergolide and/or permeation enhancer from device 40. The outer edges of backing member 14 overlay

the edges of reservoir 12 and are joined along the perimeter with the outer edges of the rate-controlling membrane 28 in a fluid-tight arrangement. This sealed reservoir may be effected by pressure, fusion, adhesion, an adhesive applied to the edges, or other methods known in the art. In this manner,
5 reservoir 12 is contained wholly between backing member 14 and rate-controlling membrane 28. On the skin-proximal side of rate-controlling membrane 28 are an adhesive layer 16 and a strippable liner 24 which would be removed prior to application of the device 40 to the skin.

In an alternative embodiment of device 40 of FIG. 4, reservoir 12
10 contains the permeation enhancer and contains pergolide at or below saturation. The pergolide and an additional amount of permeation enhancer are present in adhesive layer 16, which acts as a separate reservoir.

The pergolide with or without the permeation enhancer can be administered to human skin or mucosa by direct application to the skin or
15 mucosa in the form of an ointment, gel, cream or lotion, for example, but are preferably administered from a skin patch or other known transdermal delivery device which contains a saturated or unsaturated formulation of the pergolide or pergolide and enhancer. The formulation may be aqueous or non-aqueous based. The formulation should be designed to deliver the
20 pergolide at the necessary fluxes. Aqueous formulations typically comprise water or water/ethanol and about 1-5 wt% of a gelling agent, an example being a hydrophilic polymer such as hydroxyethylcellulose or hydroxypropylcellulose. Typical non-aqueous gels are comprised of silicone fluid or mineral oil. Mineral oil-based gels also typically contain 1-2 wt% of a
25 gelling agent such as colloidal silicon dioxide. The suitability of a particular gel depends upon the compatibility of its constituents with the drug and the permeation-enhancing mixture, if used, in addition to any other components in the formulation.

The reservoir matrix should be compatible with pergolide, the
30 permeation enhancer, and any carrier therefor. The term "matrix" as used herein refers to a well-mixed composite of ingredients fixed into shape.

When using an aqueous-based formulation, the reservoir matrix is preferably a hydrophilic polymer, eg, a hydrogel.

When using a non-aqueous based formulation, the reservoir matrix is preferably composed of a hydrophobic polymer. Suitable polymeric matrices
5 are well known in the transdermal drug delivery art, and examples are listed in the above-named patents previously incorporated herein by reference. A typical laminated system would consist essentially of a polymeric membrane and/or matrix such as ethylene vinyl acetate (EVA) copolymers, such as those described in US Pat. No. 4,144,317, preferably having a vinyl acetate
10 (VA) content in the range of from about 9% up to about 60% and more preferably about 9% to 40% VA. Polyisobutylene/oil polymers containing from 4-25% high molecular weight polyisobutylene and 20-81% low molecular weight polyisobutylene with the balance being an oil such as mineral oil or polybutene may also be used as the matrix material.

15 The amount of pergolide present in the therapeutic device and required to achieve an effective therapeutic result depends on many factors, such as the minimum necessary dosage of the pergolide for the particular indication being treated; the solubility and permeability of the matrix, taking into account the presence of a permeation enhancer, of the adhesive layer
20 and of the rate-controlling membrane, if present; and the period of time for which the device will be fixed to the skin. The minimum amount of pergolide is determined by the requirement that sufficient quantities of pergolide must be present in the device to maintain the desired rate of release over the given period of application. The maximum amount for safety purposes is
25 determined by the requirement that the quantity of pergolide present cannot exceed a rate of release that reaches toxic levels.

The pergolide is normally present in the matrix or carrier at a concentration in excess of saturation, the amount of excess being a function of the desired length of the delivery period of the system. Pergolide may,
30 however, be present at a level below saturation without departing from this invention as long as it is continuously administered to the skin or mucosal site

at a therapeutic rate and for a period of time sufficient to deliver a therapeutically effective amount of pergolide that provides the desired therapeutic result.

The permeation enhancer useful in the present invention is selected
5 from those compounds which are compatible with pergolide and which provide enhanced skin permeation to the drug when it is administered together with the drug to the skin of a user. Additionally, the permeation enhancer must not adversely interact with the adhesive of the in-line contact adhesive layer if one is present. Such permeation enhancers can be
10 selected from, but are not limited to, C₂₋₄ alcohols such as ethanol and isopropanol, polyethylene glycol monolaurate, laureth-4, lactate esters such as lauryl lactate, dimethyl lauramide, LDEA, caproyl lactic acid, esters of fatty acids having from about 10 to about 20 carbon atoms such as methyl laurate, and monoglycerides or mixtures of monoglycerides of fatty acids alone, or in
15 combinations with each other.

A preferred permeation enhancer according to this invention comprises glycerol monolaurate and methyl laurate. Methyl laurate has been found to be particularly desirable as it is obtainable at a high degree of purity, thus providing a purer and better defined permeation enhancer and a system
20 which is more readily characterized. Additionally, the inventors have found that the addition of mineral oil to the pergolide reservoir comprising glycerol monolaurate and methyl laurate provides a surprising and synergistic effect to the pergolide skin flux. In a particularly preferred embodiment, the ratio of glycerol monolaurate / methyl laurate / mineral oil is .75 / 1.0 / .50. As used
25 herein, "mineral oil" refers to a mixture of liquid hydrocarbons of petroleum. Polybutene may also be substituted for the mineral oil.

Typically, monoglycerides have been available as a mixture of monoglycerides of fatty acids with one monoglyceride being the principal component, from which component the mixture derives its name. For
30 example, one commercial monoglyceride is Emerest 2421 glycerol monooleate (Emery Division, Quantum Chemical Corp.), which is a mixture of

glycerol oleates with a glycerol monooleate content of 58% and a total monoesters content of 58%. Other examples of commercial monoglycerides are Myverol 1899K glycerol monooleate (Eastman Chemical Products) which has a glycerol monooleate content of 61% and a total monoesters content of 93%, and Myverol 1892K glycerol monolinoleate which has a glycerol monolinoleate content of 68% and a minimum total monoesters content of 90%. The monoesters are chosen from those with from 10 to 20 carbon atoms. The fatty acids may be saturated or unsaturated and include, for example, lauric acid, myristic acid, stearic acid, oleic acid, linoleic acid and palmitic acid. Monoglyceride permeation enhancers include glycerol monooleate, glycerol monolaurate and glycerol monolinoleate, for example. In a presently preferred embodiment of this invention, the permeation enhancer comprises glycerol monolaurate as the monoglyceride.

It has been seen that glycerol monooleate having a total monoesters content of less than about 65% interacts adversely with known adhesive materials to such an extent that the adhesive cannot function to maintain a delivery device on the skin. Therefore, when an in-line adhesive is present as a part of the device of the invention so that a permeation enhancer must pass through the adhesive, and when glycerol monooleate is utilized as the permeation enhancer, the glycerol monooleate must have a total monoesters content of at least 65%.

The permeation-enhancing mixture is dispersed through the matrix or carrier, preferably at a concentration sufficient to provide permeation-enhancing amounts of enhancer in the reservoir throughout the anticipated administration period. Where there is an additional, separate permeation enhancer matrix layer as well, as in FIGS. 3 and 4, the permeation enhancer normally is present in the separate reservoir in excess of saturation.

In addition to the pergolide and permeation enhancer, the matrix or carrier may also contain dyes, pigments, inert fillers, excipients and other conventional components of pharmaceutical products or transdermal devices known to the art.

Because of the wide variation in skin permeability from individual to individual and from site to site on the same body, it may be preferable that the pergolide, with or without a permeation enhancer, be administered from a rate-controlled transdermal delivery device. Rate control can be obtained
5 either through a rate-controlling membrane or adhesive or both as well as through the other means.

A certain amount of pergolide will bind reversibly to the skin, and it is accordingly preferred that the skin-contacting layer of the device include this amount of pergolide as a loading dose.

10 The surface area of the device of this invention can vary from about 1-200 cm². A typical device, however, will have a surface area within the range of about 1-50 cm², preferably about 20 cm².

The devices of this invention can be designed to effectively deliver pergolide for an extended time period of from several hours up to 7 days or
15 longer. Seven days is generally the maximum time limit for application of a single device because the adverse effect of occlusion of a skin site increases with time and the normal cycle of sloughing and replacement of the skin cells occurs in about 7 days.

Preferably, the transdermal drug delivery device contains sufficient
20 amounts of a permeation enhancer as described above and pergolide, in combination, to provide systemic administration of pergolide through the skin at a therapeutically effective rate during the administration period in order to provide therapeutic blood plasma levels.

Preferably, a device for the transdermal administration of pergolide, at
25 a therapeutically effective rate, comprises:

- (a) a reservoir comprising:
 - (i) 1-15% by weight pergolide,
 - (ii) 10-70% by weight of a permeation enhancer,
 - (iii) 35 to 85% by weight ethylene vinyl acetate having a vinyl
30 acetate content of 9-60%;
- (b) a backing behind the skin-distal surface of the reservoir; and

(c) means for maintaining the reservoir in drug- and permeation enhancer- transmitting relation with the skin.

More preferably, a device for the transdermal administration of pergolide, at a therapeutically effective rate, comprises:

- 5 (a) a reservoir comprising:
- (i) 1 to 15% by weight of a pharmaceutically acceptable salt of pergolide,
 - (ii) 10 to 70% by weight of a permeation enhancer,
 - (iii) 45 to 85% by weight ethylene vinyl acetate having a vinyl
10 acetate content of 9-40%;
- (b) a backing behind the skin-distal surface of the reservoir; and
- (c) means for maintaining the reservoir in drug- and permeation enhancer -transmitting relation with the skin.

Most preferably, a device for the transdermal administration of
15 pergolide, at a therapeutically effective rate, comprises:

- (a) a reservoir comprising:
- (i) 1 to 15% by weight pergolide mesylate,
 - (ii) 1-35% by weight glycerol monolaurate,
 - (iii) 1-35% by weight methyl laurate,
 - (iv) 1-35% by weight mineral oil,
20 (v) 45 to 80% by weight ethylene vinyl acetate having a vinyl acetate content of 30-40%;
- (b) a backing behind the skin-distal surface of the reservoir; and
- (c) means for maintaining the reservoir in drug- and permeation
25 enhancer- transmitting relation with the skin.

The backing may be a breathable or occlusive material such as polyethylene, polyurethane, polyester or ethylene vinyl acetate films. If mineral oil is used, a polyester backing is preferred. If an ethylene vinyl acetate is employed as the backing, preferably, it has an acetate content of
30 33% or 40%.

The means for maintaining the reservoir in drug and permeation enhancer transmitting relation with the skin are preferably a polyisobutylene adhesive, as described in the Examples that follow. A further embodiment of the invention is directed to including in the adhesive a small percentage, eg,
5 from about 1.0 to about 5 wt% of pergolide.

The aforementioned patents describe a wide variety of materials which can be used for fabricating various layers or components of the transdermal pergolide delivery systems according to this invention. This invention, therefore, contemplates the use of other materials other than those
10 specifically disclosed herein including those which may become hereafter known to the artist capable of forming the necessary functions.

The invention is also directed to a method of continuously administering pergolide to a individual at a therapeutically effective rate during an administration period in order to provide substantially constant
15 therapeutic blood plasma levels of pergolide in an individual.

Another method of the present invention is directed to a method for the transdermal coadministration of pergolide at a therapeutically effective rate together with a skin permeation-enhancing amount of a permeation enhancer in order to achieve and maintain therapeutic blood plasma levels of pergolide
20 in an individual, comprising:

- (a) coadministering to a body surface or membrane, pergolide; and
- (b) a permeation enhancer, wherein pergolide is delivered at a therapeutically effective rate during the administration period in order to achieve and maintain therapeutic blood plasma levels of pergolide in an
25 individual. The pergolide and permeation enhancer may be administered to the body surface or membrane by means of the devices and compositions described above.

A preferred embodiment of the present invention comprises a method of treating Parkinson's Disease. To be useful in treating Parkinson's
30 Disease, pergolide should be present above plasma concentrations of about 100 pg/mL, preferably at concentrations above about 300 pg/mL and most

preferably at concentrations of about 1000 pg/mL. To achieve this result, pergolide is delivered at an estimated therapeutic rate of at least about 100 μg per hour, but typically of at least 125 $\mu\text{g/hr}$, and more typically at about 150 $\mu\text{g/hr}$, for the treatment period, usually about 12 hours to 7 days. For example, a 20 cm^2 system would require a pergolide flux through skin of 7.5 $\mu\text{g/cm}^2 \cdot \text{hr}$ in order to achieve the desired therapeutic rate of 150 $\mu\text{g/hr}$. Alternatively, three 25 cm^2 systems would require a pergolide flux of about 2 $\mu\text{g/cm}^2 \cdot \text{hr}$ to achieve the administration rate of 150 $\mu\text{g/hr}$. Such regimens would deliver approximately 3.6 mg of pergolide over a 24 hour period.

The length of time of pergolide presence and the total amount of pergolide in the plasma can be changed following the teachings of this invention to provide different treatment regimens. Thus, they can be controlled by the amount of time during which exogenous pergolide is delivered transdermally to an individual or animal.

Having thus generally described our invention, the following specific examples describe preferred embodiments thereof but are not intended to limit the invention in any manner.

EXAMPLE 1

Several test samples were made to measure the flux of pergolide base and mesylate through human cadaver epidermis from donor vehicles containing the pergolide base or mesylate mixed with water alone, water and ethanol, or mineral oil alone as shown in Table 1. Transdermal fluxes were obtained using human epidermis at 35°C in standard diffusion cells. Figures 5 and 6 graphically depict the results. As seen in these Figures, the average baseline skin flux of pergolide base without any permeation enhancer is approximately 0.3 $\mu\text{g/cm}^2 \cdot \text{hr}$ over a fifty two hour period, while the average baseline skin flux of pergolide mesylate without permeation enhancers is approximately 1.1 $\mu\text{g/cm}^2 \cdot \text{hr}$ over the same time period.

TABLE 1**Aqueous and Non-Aqueous Donor Solutions (weight percent)**

Pergolide (form and content)	H₂O	EtOH	Mineral Oil
Base 2	98	0	0
Base 5	45	55	0
Base 5	70	30	0
Base 2	0	0	98
Mesylate 2	98	0	0
Mesylate 5	45	55	0
Mesylate 5	70	30	0
Mesylate 2	0	0	98

EXAMPLE 2

5

The drug/permeation enhancer reservoirs were prepared by mixing ethylene vinyl acetate having a vinyl acetate content of 40 percent ("EVA 40", USI Chemicals, Illinois), with pergolide base or mesylate, GML (Grindsted Products, Braband, Denmark) or LDEA (Lonza, Inc.), and mineral oil (Witco Corp.). The mixture was then dissolved in tetrahydrofuran. After blending, the mixture was hand cast and dried to a 5 mil. thick film. The composition of the drug reservoirs is shown in Table 2.

10

TABLE 2
Drug/Permeation Enhancer Reservoir Composition (weight percent)

Pergolide (form and content)	LDEA	Glycerol Monolaurate	Mineral Oil	EVA 40
Base 10	15	0	15	60
Base 10	0	20	14	56
Mesylate 10	15	0	15	60
Mesylate 10	0	20	14	56

5 The film was then laminated to a pigmented medium density polyethylene/aluminum foil/PET/EVA (Medpar®) backing on one side and an acrylate contact adhesive on the opposite side (3M, St. Paul, MN). The laminate was then punched down to an area of 1.6 cm².

10 Circular pieces of human epidermis were placed with stratum corneum facing up. The release liner of the laminate was removed and the system was centered over the stratum corneum side of the epidermis. The edges of epidermis were then folded around the system. This assembly was then mounted on a Teflon rod. A known volume of receptor solution was then placed in a test tube and was equilibrated at 35°C. The Teflon rod with
15 system and epidermis attached was then placed in a water bath at 35°C. Mixing was accomplished by attachment to a motor which caused constant vertical mixing.

 At given time intervals, the entire receptor solution was removed from the test tubes and replaced with an equal volume of fresh receptor solutions
20 previously equilibrated at 35°C. The receptor solutions are stored in capped vials at room temperature until assayed for pergolide content by HPLC. From the drug concentration and the volume of the receptor solutions, the area of permeation and the time interval, the flux of the drug through the epidermis

was calculated as follows: (drug concentration X volume of receptor)/(area x time) = flux ($\mu\text{g}/\text{cm}^2 \cdot \text{hr}$). The cumulative release of the pergolide base or mesylate is shown in Figures 7 and 8.

5

EXAMPLE 3

The drug/permeation enhancer reservoirs were prepared according to Example 2. The film was then laminated to Medpar backing on one side and a polyisobutylene adhesive containing 2.5% by weight of pergolide on the other. The adhesive was prepared by dissolving 19.8% 1.2M polyisobutylene, 24.7% 35K polyisobutylene and 55.5% light mineral oil in heptane. The 2.5% pergolide was added and the entire mixture was cast to a dry thickness of 2 mils. The film was then cut into circles using a stainless steel punch with an area of 1.6 cm^2 .

15 The epidermis was separated from the dermis of the skin donor after immersion in 60°C water for 60 seconds. Discs (7/8-inch diameter) were cut from the epidermis, and the discs were kept at 4°C in a hydrated state until they were used.

For each device tested, the release liner was removed and the drug-releasing surface was placed against the stratum corneum side of a disc of epidermis which had been blotted dry just prior to use. The excess epidermis was wrapped around the device so that none of the device edge was exposed to the receptor solution. The device covered with epidermis was attached to the flat side of the Teflon holder of a release rate rod using nylon netting and nickel wire. The rods were reciprocated in a fixed volume of receptor solution. The entire receptor solution was changed at each sampling time. The temperature of the receptor solution in the water bath was maintained at 35°C . The cumulative release of the pergolide base or pergolide mesylate is shown in Figures 9 and 10.

30

EXAMPLE 4

The drug/permeation enhancer reservoirs were prepared by mixing pergolide base or mesylate, EtOH, GML, and caproyl lactic acid (CLA)
5 (R.I.T.A. Corp., Woodstock, IL) as shown in Table 3.

TABLE 3

Drug/Permeation Enhancer Reservoir Composition (weight percent)

Pergolide (form and content)	EtOH	GML	CLA
Base 10	80	10	0
Base 10	90	0	0
Base 10	85	0	5
Mesylate 10	80	10	0
Mesylate 10	90	0	0
Mesylate 10	85	0	5

An additional 3% by weight hydroxypropylcellulose and water were added
10 and the mixture was placed in a suitable container and gelled. A desired quantity of the mixture was then applied on the surface of a previously formed trilaminate consisting of a 2 mil thick ethylene vinyl acetate film having a vinyl acetate content of 18% for use as a rate controlling membrane; a polyisobutylene contact adhesive containing 2.5% by weight pergolide; and
15 a polyethylene ethylene terephthalate film coated with silicone for use as a release liner. A Medpar backing was then applied on top of the mixture and the entire system was heat sealed. The systems were then die-cut to the required sizes for use in the in vitro test method described in Examples 2 & 3. The cumulative release of the pergolide base and mesylate are shown in
20 Figures 11 and 12.

EXAMPLE 5

A desired quantity of the mixture as formed in Example 4 was then applied on the surface of a previously formed trilaminate consisting of a 2 mil thick ethylene vinyl acetate film having a vinyl acetate content of 28% for use as a rate controlling membrane; a polyisobutylene contact adhesive containing 2.5% by weight pergolide; and a polyethylene ethylene terephthalate film coated with fluorocarbon for use as a release liner. A Medpar backing was then applied on top of the mixture and the entire system was heat sealed. The systems were then die-cut to the required sizes for use in the in vitro test method described in Examples 2-4. The skin flux of the base and mesylate are shown in Figures 13 and 14.

EXAMPLE 6

The effect of mineral oil on the permeation enhancement of GML with various cosolvents was studied. The drug/permeation enhancer reservoirs were prepared by mixing ethylene vinyl acetate having a vinyl acetate content of 40 percent ("EVA 40", USI Chemicals, Illinois), pergolide mesylate, GML, and a cosolvent selected from Laureth-4 (L-4) (Heterene Chemical Co., Inc., Paterson, N.J.), methyl laurate (Sigma), lauryl lactate (ISP Van Dyk Inc., Belleville, NJ) and dodecyl acetate (Penta). The mixture was then dissolved in tetrahydrofuran. After blending, the mixture was hand cast and dried to a 5 mil. thick film. Various compositions for each cosolvent without mineral oil were compared with a control composition comprising mesylate/GML/laureth-4/EVA 10/20/12/58. The compositions of the drug reservoirs is shown in Tables 4a-4c.

TABLE 4a

Drug/Permeation Enhancer Reservoir Composition (weight percent)

Pergolide mesylate	GML	Methyl Laurate	EVA 40
5	17	3	75
5	15	5	75
5	10	10	75

TABLE 4b

5 Drug/Permeation Enhancer Reservoir Composition (weight percent)

Pergolide mesylate	GML	Dodecyl Acetate	EVA 40
5	17	3	75
5	15	5	75
5	10	10	75

TABLE 4c

Drug/Permeation Enhancer Reservoir Composition (weight percent)

Pergolide mesylate	GML	Lauryl Lactate	EVA 40
5	17	3	75
5	15	5	75
5	10	10	75

10

The film was then laminated to a Medpar backing on one side. The systems were then die-cut to the required sizes for use in the in vitro test method described in Examples 2-4. The skin flux of pergolide mesylate is shown in Figures 15 - 17.

15

Compositions comprising 5% pergolide mesylate, 10% GML, 10% of one of the lauryl lactate, methyl laurate, or dodecyl acetate cosolvents, 15% mineral oil, and 60% EVA 40 were then prepared and used in the in vitro skin flux experiment described above and compared to the pergolide

mesylate/GML/laureth 4/EVA 40 10/20/12/58 control. The result of the in vitro skin flux experiment is shown in Figure 18. As seen in Figure 18, the mineral oil had the most substantial effect on the composition comprising GML and methyl laurate.

5 Having thus generally described our invention and described certain specific embodiments thereof, including the embodiments that applicants consider the best mode of practicing their invention, it should be readily apparent that various modifications to the invention may be made by workers skilled in the art without departing from the scope of this invention which is
10 limited only by the following claims.

What is claimed is:

1. A composition of matter for delivering pergolide through a body surface or membrane at a therapeutically effective rate to an individual in
5 need of pergolide therapy by permeation through the body surface or membrane, the composition comprising an amount of pergolide or a pharmaceutically acceptable salt in a carrier effective to permit sustained release of pergolide over an administration period at a flux of at least about 1 $\mu\text{g}/\text{cm}^2 \cdot \text{hr}$ in order to achieve therapeutic blood plasma levels of pergolide.
10
2. A composition according to claim 1 further comprising a permeation enhancer selected from the group consisting of lauramide diethanolamine, glycerol monolaurate, glycerol monooleate, methyl laurate, lactate esters, caproyl lactic acid, ethanol, and polyethylene glycol-4 lauryl
15 ether, alone or in combinations of one or more, in an amount sufficient to substantially increase the permeability of the body surface or membrane to pergolide.
3. A composition according to claim 2 wherein the pergolide is
20 provided as a pharmaceutically acceptable salt.
4. A composition according to claim 3 wherein the pharmaceutically acceptable salt is pergolide mesylate.
- 25 5. A composition according to claim 4 wherein the permeation enhancer is a mixture comprising glycerol monolaurate and methyl laurate.
6. A composition according to claim 5 further comprising mineral
oil.

7. A device for the transdermal administration of pergolide at a therapeutically effective rate to an individual in need of pergolide therapy, comprising:

- (a) a reservoir comprising an amount of pergolide;
- 5 (b) a backing behind the body contacting-distal surface of the reservoir; and
- (c) means for maintaining the reservoir in drug transmitting relation with a body surface or membrane wherein pergolide is provided in an amount effective to permit sustained release of pergolide during an
- 10 administration period at a flux of at least about $1 \mu\text{g}/\text{cm}^2 \cdot \text{hr}$ in order to achieve therapeutic blood plasma levels of pergolide.

8. A device according to claim 7 wherein the reservoir further comprises a permeation enhancer selected from the group consisting of

15 lauramide diethanolamine, glycerol monolaurate, glycerol monooleate, methyl laurate, lactate esters, caproyl lactic acid, ethanol, and polyethylene glycol-4 lauryl ether, alone or in combinations of one or more, in an amount sufficient to substantially increase the permeability of the body surface or membrane to the pergolide.

20

9. A device according to claim 8 wherein the pergolide is provided as a pharmaceutically acceptable salt.

10. A device according to claim 9 wherein the pharmaceutically

25 acceptable salt is pergolide mesylate.

11. A device according to claim 10 wherein the means for maintaining the reservoir in relation with the skin comprises an in-line adhesive layer on the body contacting-proximal surface of the reservoir.

30

12. A device according to claim 11 wherein the in-line adhesive is a polyisobutylene adhesive.

13. A device according to claim 10 wherein the permeation
5 enhancer comprises glycerol monolaurate and methyl laurate.

14. A device according to claim 13 wherein the reservoir further comprises mineral oil.

10 15. A device for the transdermal administration of pergolide at a therapeutically effective rate to an individual in need of pergolide therapy, comprising:

(a) a first reservoir comprising an amount of pergolide and a permeation-enhancing amount of a permeation enhancer;

15 (b) a second reservoir comprising an excess of the permeation enhancer and pergolide at or below saturation when in equilibrium with the first reservoir;

(c) a rate-controlling membrane between the first reservoir and the second reservoir;

20 (d) a backing behind the body contacting-distal surface of the second reservoir; and

(e) means for maintaining the first and second reservoirs in drug-and permeation enhancer-transmitting relation with a body surface or membrane wherein pergolide is provided in an amount effective to permit
25 sustained release of pergolide during an administration period at a flux of at least about $1 \mu\text{g}/\text{cm}^2 \cdot \text{hr}$ in order to achieve therapeutic blood plasma levels of pergolide.

16. A device according to claim 15 wherein the permeation
30 enhancer is selected from the group consisting of lauramide diethanolamine, glycerol monolaurate, glycerol monooleate, methyl laurate, lactate esters,

caproyl lactic acid, ethanol, and polyethylene glycol-4 lauryl ether, alone or in combinations of one or more, in an amount sufficient to substantially increase the permeability of the body surface or membrane to pergolide.

5 17. A device according to claim 16 wherein the pergolide is provided as a pharmaceutically acceptable salt.

 18. A device according to claim 17 wherein the pharmaceutically acceptable salt is pergolide mesylate.

10

 19. A device according to claim 18 wherein the permeation enhancer comprises glycerol monolaurate and methyl laurate.

 20. A device according to claim 19 wherein the reservoir further
15 comprises mineral oil.

 21. A device according to claim 20 wherein the means for maintaining the reservoir in relation with the skin comprises an in-line adhesive layer on the body contacting-proximal surface of the reservoir.

20

 22. A device according to claim 21 wherein the in-line adhesive is a polyisobutylene adhesive.

 23. A device for the transdermal administration of pergolide at a
25 therapeutically effective rate to an individual in need of pergolide therapy, comprising:

 (a) a reservoir comprising:

 (i) 1 to 15 wt% of pergolide;

 (ii) 10 to 70 wt% of permeation enhancer selected

30 from the group consisting of lauramide diethanolamine, glycerol monolaurate, glycerol monooleate, methyl laurate, lactate esters, caproyl lactic acid,

ethanol, and polyethylene glycol-4 lauryl ether, alone or in combinations of one or more;

(iii) 35 to 85 wt% of ethylene vinyl acetate copolymer;

(b) a backing behind the body contacting-distal surface of
5 the reservoir; and

(c) means for maintaining the reservoir in drug-and permeation enhancer-transmitting relation with a body surface or membrane, wherein pergolide is provided in an amount effective to permit sustained release of pergolide during an administration period at a flux of at least about
10 $1 \mu\text{g}/\text{cm}^2 \cdot \text{hr}$ in order to achieve therapeutic blood plasma levels of pergolide.

24. A device according to claim 23 wherein the pergolide is provided as a pharmaceutically acceptable salt.

15 25. A device according to claim 24 wherein the pharmaceutically acceptable salt is pergolide mesylate.

26. A device according to claim 25 wherein the permeation enhancer comprises 1-35 wt% glycerol monolaurate and 1-35 wt% methyl
20 laurate.

27. A device according to claim 26 wherein the reservoir further comprises 1-35 wt% mineral oil.

25 28. A device according to claim 27 wherein the reservoir comprises a ratio of glycerol monolaurate / methyl laurate / mineral oil of .75 / 1.0 / .50.

29. A method for the transdermal administration of pergolide to an individual in need of pergolide therapy by permeation through a body surface
30 or membrane at a therapeutically effective rate, which method comprises administering pergolide to a body surface or membrane in a carrier effective

to permit sustained release of pergolide during an administration period at a flux of at least about $1 \mu\text{g}/\text{cm}^2 \cdot \text{hr}$ in order to achieve therapeutic blood plasma levels of pergolide.

5 30. A method according to claim 29 further comprising simultaneously coadministering a permeation-enhancing amount of a permeation enhancer selected from the group consisting of lauramide diethanolamine, glycerol monolaurate, glycerol monooleate, methyl laurate, lactate esters, caproyl lactic acid, ethanol, and polyethylene glycol-4 lauryl
10 ether, alone or in combinations of one or more, in an amount sufficient to substantially increase the permeability of the body surface or membrane to the pergolide.

 31. A method according to claim 30 wherein pergolide is provided
15 as a pharmaceutically acceptable salt.

 32. A method according to claim 31 wherein the pharmaceutically acceptable salt is pergolide mesylate.





20 33. A method according to claim 32 wherein the permeation enhancer comprises glycerol monolaurate and methyl laurate.

 34. A method according to claim 33 wherein the carrier further comprises mineral oil.
25

 35. A method for treating Parkinson's Disease comprising administering pergolide or a pharmaceutically acceptable salt to a body surface or membrane in a carrier effective to permit sustained release of pergolide during an administration period at a flux of at least about $1 \mu\text{g}/\text{cm}^2 \cdot \text{hr}$ in order to achieve therapeutic blood plasma levels of pergolide.
30


36. A method according to claim 35 further comprising simultaneously coadministering a permeation-enhancing amount of a permeation enhancer selected from the group consisting of lauramide diethanolamine, glycerol monolaurate, glycerol monooleate, methyl laurate, lactate esters, caproyl lactic acid, ethanol, and polyethylene glycol-4 lauryl ether, alone or in combinations of one or more, in an amount sufficient to substantially increase the permeability of the body surface or membrane to the pergolide.

1 / 15


**FIG. 1****FIG. 2****FIG. 3****FIG. 4**

2 / 15


FIG. 5




3 / 15





4 / 15






6 / 15






* Expressed as Base

8 / 15





9 / 15



* Expressed as Base

10 / 15







* Expressed as Base

12 / 15

FIG. 15



13 / 15



14 / 15





FIG. 18



INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/09692

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/48 A61K9/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE,A,42 40 798 (POLI INDUSTRIA CHIMICA S.P.A.,IT) 9 June 1993 cited in the application see the whole document ---	1-36
A	EP,A,0 458 640 (ELI LILLY AND COMPANY,U.S.A.) 27 November 1991 cited in the application see the whole document ---	1-36
A	WO,A,91 00746 (CYGNUS RESEARCH CORPORATION,U.S.A.) 24 January 1991 cited in the application see the whole document -----	1-36

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

- * "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- * "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- * "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- * "&" document member of the same patent family

Date of the actual completion of the international search

28 October 1996

Date of mailing of the international search report

07.11.96

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Scarponi, U

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/09692

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 35-36
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/09692

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A-4240798	09-06-93	IT-B- 1252163	05-06-95

EP-A-458640	27-11-91	US-A- 5063234	05-11-91
		AU-B- 631991	10-12-92
		AU-A- 7730391	28-11-91
		CA-A- 2042960	26-11-91
		JP-A- 4235129	24-08-92

WO-A-9100746	24-01-91	US-A- 5252335	12-10-93
		AU-B- 648031	14-04-94
		AU-A- 6034090	06-02-91
		EP-A- 0483213	06-05-92
		JP-T- 4506958	03-12-92
		US-A- 5229129	20-07-93
